



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GERMANY.

Report from Berlin—Death rate of Berlin and other cities.

Consul-General Mason reports, July 25, as follows:

The death rate of Berlin for the week ended July 8 was higher than it has been for 10 months past, amounting, calculated on the year, to 19.1 per thousand of the population, thus being also considerably higher than the rate for the same week of last year, in which it amounted only to 14.5 per thousand. Almost two-thirds of the large towns and cities of Germany, however, showed still more unfavorable health conditions than Berlin. The following cities had a higher rate of mortality than Berlin, namely: Hamburg, Bremen, Frankfort on the Main, Munich, Leipzig, Dresden, Rixdorf (with 38 per thousand), Königsberg, Breslau, Halle, Magdeburg, Nuremberg, Stuttgart, Karlsruhe, Strassburg, and also Vienna. The following cities had a lower death rate than Berlin, viz: Schöneberg (with 13.2), Charlottenburg (with 15.4), Brunswick, Hanover, Essen, Altona, Kiel, as well as London and Paris. The infant death rate rose from 6.1 per year and thousand to 7.9. There were registered 95 deaths from phthisis pulmonalis; 87 deaths from acute diseases of the respiratory organs; 45 deaths from cancer; 8 deaths from measles; 6 deaths from scarlet fever; 4 deaths from diphtheria; 2 deaths from influenza; and, finally, 27 persons died by violence.

GUATEMALA.

Report from Livingston, fruit port—Yellow fever.

Acting Assistant Surgeon Peters reports as follows: Week ended July 22, 1905. Present officially estimated population, 3,500; 4 deaths, including 2 from yellow fever; prevailing diseases, malarial and yellow fever; general sanitary condition of this port and the surrounding country during the week, infected.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
July 18	Spero.....	18
18	Anselm.....	40
20	Anselm.....	40

The steamship *Anselm* cleared from Puerto Barrios and Livingston.

Yellow fever reported present at Zacapa.

PUERTO BARRIOS, August 3, 1905.

WYMAN, *Washington*:

It is reported that there have been 9 deaths from yellow fever, Zacapa, on railroad above Barrios.

PETERS.